

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problems Mailbox.**

# Exhibit K

# Leucocyte Typing IV

## White Cell Differentiation Antigens

---

Edited by

W. Knapp

*Institute for Immunology, University of Vienna*

B. Dörken

*University of Heidelberg*

E. P. Rieber

*University of Munich*

H. Stein

*Freie Universität Berlin*

W. R. Gilks

*MRC Biostatistics Unit, Cambridge*

R. E. Schmidt

*Medizinische Hochschule, Hannover*

A. E. G. Kr. von dem Borne

*CLB Netherlands Red Cross and University of  
Amsterdam, Amsterdam*

Oxford New York Tokyo

OXFORD UNIVERSITY PRESS

1989

Oxford University Press, Walton Street, Oxford OX2 6DP  
 Oxford New York Toronto  
 Delhi Bombay Calcutta Madras Karachi  
 Petaling Jaya Singapore Hong Kong Tokyo  
 Nairobi Dar es Salaam Cape Town  
 Melbourne Auckland  
 and associated companies in  
 Berlin Ibadan

Oxford is a trade mark of Oxford University Press

Published in the United States  
 by Oxford University Press, New York

© The Organizing Committee of the Fourth International  
 Conference on Human Leucocyte Differentiation Antigens and Oxford University Press, 1989

All rights reserved. No part of this publication may be reproduced,  
 stored in a retrieval system, or transmitted, in any form or by any means,  
 electronic, mechanical, photocopying, recording, or otherwise, without  
 the prior permission of Oxford University Press

British Library Cataloguing in Publication Data

Leucocyte typing IV.

1. Man. Blood. White cells. Antigens

I. Knapp, W. (Walter)

612'.112

ISBN 0-19-261867-9

Library of Congress Cataloging in Publication Data

International Workshop and Conference on Human Leucocyte

Differentiation Antigens (4th : 1989 : Vienna, Austria)

Leucocyte typing IV : white cell differentiation antigens/edited

by W. Knapp . . . [et al.].

"Fourth International Workshop and Conference on Human Leucocyte  
 Differentiation Antigens, Vienna, 21-25 February 1989"—P. facing t.p.

Includes bibliographical references.

1. HLA histocompatibility antigens—Classification—Congresses.

2. Histocompatibility testing—Congresses I. Knapp, W. (Walter)

II. Title. III. Title : Leucocyte typing 4.

[DNLM : 1. Antibodies, Monoclonal—classification—congresses.

2. Antigens, Differentiation—congresses. 3. Leukocytes—

classification—congresses. 4. Leukocytes—immunology—congresses.

WH 200 I616L 1989]

QR184.32.I58 1989 616.07'92—dc20 89-23017 CIP

ISBN 0-19-261867-9

Typeset by Cotswold Typesetting, Gloucester

Printed in Great Britain by

Butler & Tanner Ltd

Frome, Somerset

## 1088 CD Guide

## Appendix A

## CD51 (cont.)

25 kDa) is non-covalently associated with the VNR  $\beta$ 3 chain, which is identical to GPIIb (CD61 molecule).

**Molecular mass** 125 kDa (large subunit VNR  $\alpha$ -chain); 25 kDa (small subunit VNR  $\alpha$ -chain).

**Main cellular reactivity** Platelets (weak).

**Other reactive cells** Immunohistology: endothelial cells, smooth muscle cells, fibroblasts, osteoclasts, melanoma cells.

**Functions** VNR mediates cell adhesion to arg-gly-arg-containing sequences in vitronectin, von Willebrand factor, fibrinogen, thrombospondin.

**Selected references**

- \*Fitzgerald, L. A., Poncz, M., Steiner, B., Rall, S. C., Bennett, J. S., and Phillips, D. R. *Biochemistry* 26, 8158 (1987).  
 \*Suzuki, S., Argraves, W. S., Arai, H., Langino, L. R., Piersbacher, M. D., and Ruoslahti, E. *J. Biol. Chem.* 262, 14080 (1987).  
 Lam, S. C. T., Plow, E. F., D'Souza, S. E., Cheresh, D. A., Frelinger III, A. L., and Ginsberg, M. H. *J. Biol. Chem.* 264, 3742 (1989).

## CDw52

**Antibodies assigned at Fourth International Workshop** 097; YTH66.9; YTH361.10; YTH34.5.

**Other names** Campath-1

**Molecular structure** O-linked carbohydrate as antigenic determinant.

**Molecular mass** 21-28 kDa (?).

**Main cellular reactivity** Leucocytes (neutrophils weak, red cells and platelets negative).

**Other reactive cells** Flow cytometry: eosinophils.

**Functions** One antibody (YTH361.10) of this series is mitogenic.

**Clinical research applications** The Campath-1 antibody has been used for prevention of graft vs. host disease in allogeneic bone marrow transplantation.

**Selected references**

- Hale, G., Bright, S., Chumbley, G., Hoang, T., Metcalf, D., Munro, A., and Waldman, H. *Blood* 62, 873 (1983).  
 Hale, G., Cobbold, S., Waldmann, H., Easter, G., Matejschuk, P., and Coombs, R. A. *J. Immunol. Meth.* 103, 59 (1987).  
 Hale, G., Xia, M., and Waldman, H., this volume [N12.1], (*Leucocyte typing IV* (ed. W. Knapp, B. Dörken, W. R. Gilks, E. P. Rieber, R. E. Schmidt, H. Stein, and A. E. G. Kr. von dem Borne), Oxford University Press (1989)).

## CD53

**Antibodies assigned at Fourth International Workshop** HI29; HI36; MEM-53; HD77.

**Molecular structure** Single-chain glycoprotein.

**Molecular mass** 32-40 kDa.

**Main cellular reactivity** Broad pan-leucocyte including normal and neoplastic plasma cells.

**Other reactive cells** Immunohistology: Bone marrow cells, including osteoblasts and osteoclasts.

**Functions** Stimulation of the oxidative burst.

**Clinical research applications** CD53 mAb may be a useful tool for the discrimination of haematopoietic neoplasms from sarcomas and melanomas.

**Selected references**

- Stevanova et al., this volume [N8.1], (*Leucocyte typing IV* (ed. W. Knapp, B. Dörken, W. R. Gilks, E. P. Rieber, R. E. Schmidt, H. Stein, A. E. G. Kr. von dem Borne), Oxford University Press (1990)).  
 Hadam, M. K., this volume [N13], (*Leucocyte typing IV* (ed. W. Knapp, B. Dörken, W. R. Gilks, E. P. Rieber, R. E. Schmidt, H. Stein, and A. E. G. Kr. von dem Borne), Oxford University Press (1989)).

## CD54

**Antibodies assigned at Fourth International Workshop** RR1/1; LB-2; 7P7; 8P5; WEHI-CAM1; OKT27; P2B1.8; My13.

**Other names** Intercellular adhesion molecule-1 (ICAM-1).

**Molecular structure** Integral membrane glycoprotein with seven potential N-linked glycosylation sites.

**Molecular mass** 90 kDa.

**Main cellular reactivity** Endothelial cells, many cell types upon activation.

**Functions** ICAM-1 is a lymphokine inducible molecule, has shown to be a ligand for LFA-1 mediated adhesion, a rhinovirus receptor.

**Selected references**

- Dustin, M. L., Staunton, D. E., and Springer, T. A. *Immunol. Today* 9, 213 (1988).  
 Simmons, D., Makgoba, M. W., and Seed, B. *Nature* 331, 624 (1988).

## CD55

**Antibodies assigned at Fourth International Workshop** 143-30; BRIC110; BRIC128; F2B-7.2.

**Other names** Decay accelerating factor (DAF).

**Molecular structure** PI-linked single-chain glycoprotein, alternate splicing generates soluble form (?).

**Molecular mass** 70 kDa.

**Main cellular reactivity** All haematopoietic cells and many non-haematopoietic cells.

**Functions** DAF is involved in limiting complement activation on autologous tissue.

**Clinical research applications** Diagnosis of paroxysmal nocturnal haemoglobinuria (PNH).

**Selected references**

- Knosha, T., Medof, E., Silber, R., and Nussenzweig, V. *J. exp. Med.* 162, 75 (1985).  
 \*Caras, L. W., Davitz, M. A., Rhoe, L., Weddell, G., Martin, D. W., Jr., and Nussenzweig, V. *Nature* 325, 545 (1987).  
 \*Medof, M. E., Walter, E. I., Rutgers, J. L., Knowles, D. M., and Nussenzweig, V. *J. exp. Med.* 165, 838 (1987).  
 Lublin, D. M., Lisowski, M. K., Ost, T. W., Arce, M. A., Le Beau, M. M., Reberich, M. B., Lemons, R. S., Seys, T., and Atkinson, J. P. *J. exp. Med.* 168, 181 (1988).

## CD56

**Antibodies assigned at Fourth International Workshop** NKH1A; Leu19; FP2-11.14; L185; N901; T-199.

**Other names** Isoform of N-CAM, NKH1.

**Molecular structure** Heavily glycosylated 140 kDa isoform of N-CAM.

**Molecular mass** 220/135 kDa.

**Main cellular reactivity** NK-cells.

**Other reactive cells** Flow cytometry: neuroectodermal cells, some T-cell lines.

**Functions** Homotypic adhesion.

**Clinical research applications** Identification of NK-cells.

**Selected references**

- Hercend, T., Griffin, J. D., Bensussan, A., Schmidt, R. E., Edson, M. A., Brennan, A., Murray, C., Daley, J. F., Schlossman, S. F., and Ritz, J. *J. clin. Invest.* 75, 932 (1985).  
 Lanier, L. L., Li, A. M., Clavin, C. E., Loken, M. R., and Phillips, J. H. *J. Immunol.* 136, 4480 (1986).  
 \*Cunningham, B. A., Hemperly, J. J., Murray, B. A., Prediger, E. A., Backenbury, R., and Edelman, G. M. *Science* 236, 799 (1987).  
 Schmidt, R. E., Michon, J. M., Woronicz, J., Schlossman, S. F., Reinherz, E. L., and Ritz, J. *J. clin. Invest.* 79, 305 (1987).

N15.1

N16 CD56

NK- and Non-lineage antigens — papers 699

d cells

## N16 Cluster report: CD56

JÖRG SCHUBERT, LEWIS L. LANIER, and REINHOLD E. SCHMIDT

have shown  
M<sub>r</sub> 3000.  
agglutination  
of red cells,  
up antigens  
as described  
M<sub>r</sub> 70 000  
complement  
DAF) [4, 5].  
has been  
potential N-  
lation sites,  
glycosylphos-  
F and other  
bution have

A new cluster defined by six monoclonal antibodies (mAb) from five different laboratories which have binding specificity to the N-CAM/NKH1 molecule could be established. The molecule has previously been characterized on peripheral blood lymphocytes (PBL) as NKH1. It is expressed on all cells mediating non-MHC restricted cytotoxicity and is, therefore, a pan-NK cell antigen [1]. NKH1A was the first mAb generated by immunization against a human NK-clone [2]. It was found to detect the same antigen as the previously published N901 (NKH1) antibody [3]. This antigen has been extensively studied in its relation to NK-cells. It is detected on a subpopulation of approximately 10–15 per cent of human PBL [2–4]. Depletion of NKH1<sup>+</sup> cells by complement lysis completely abolishes non-MHC restricted cytotoxicity of PBMNC [2]. Moreover, it can be shown that the vast majority of NK active cells express the NKH1 molecule. FACS sorting of PBL demonstrated all NK cytotoxicity to be contained within the NKH1<sup>+</sup> population [5]. The molecule also appears to represent a late activation antigen since it can be induced on various long-term cultured interleukin-2-dependent T-cell clones [6]. These NKH1<sup>+</sup> T-cell clones were described as cytotoxic as well as non-cytotoxic.

Lanier *et al.* [7] presented data indicating that the NKH1 molecule is identical to the 140-kDa isoform of human neuronal cellular adhesion molecule N-CAM. In previous studies it has been demonstrated that the N-CAM molecule exists in at least three different isoforms [8]. A 120-kDa molecule which has a PI-linked membrane anchor, a 140 and a 160-kDa isoform have been described [8–10]. The isoforms are products of different RNA splicing from a single gene located on chromosome 11 [11–13] and have been demonstrated to play a role in neuronal homotypic cell adhesion and cell differentiation during embryogenesis [14].

Immunoprecipitation from NK-cells and from neuroblastoma using either polyvalent rabbit antisera against human N-CAM or the NKH1-specific mAb Leu-19 and subsequent deglycosylation revealed an identical band of 140 kDa [7]. In neuronal tissue three different isoforms can be detected. In contrast, NK-cells express the 140-kDa molecule only. In both cell types the molecule contains about 25–30 per cent of N-linked sugars and sialic acid.

Three mAb have been submitted to the new antibody

panel as recognizing the NKH1 antigen: N3 (Leu-19), N9 (N901), and N77 (NKH1A). As control antibody Leu-19 was utilized as a duplicate (N3=N79) in the panel. This reagent proved to be very useful in evaluating functional studies. During the Workshop pre-screening procedure three additional antibodies from the new antibody panel submitted with unknown specificities were assigned to the N-CAM/NKH1 group: N21 (FP2-11.14), N63 (T-199), and N126 (L185). On evaluating the phenotypic, biochemical, and histological data the six antibodies define a new cluster. The phenotypic pattern of the antibodies is shown in Fig. 1. From the histological data it was suggested that, in addition to N-CAM/NKH1 specificity, another reactivity might be detected in N63 (T-199). Binding specificity of the grouped antibodies was also studied biochemically. Immunoprecipitation demonstrated a single band of approximately 160 kDa. This precipitation was performed by Perussia from cultured NK-cells. Gelsberg *et al.* [N21.1] detected two bands of 220 and 135 kDa when using LGL-leukaemia cells as target (Table 1). Antibodies NKH1A and L185, both of IgM isotype, failed to precipitate specific bands. The differences in molecular weight may correspond to differences in glycosylation of the molecules on the different targets used for precipitation. The additional band of 135 kDa from the precipitation of LGL-leukaemia cells obviously represents the unglycosylated core protein.

For epitope mapping, cross-blocking studies were performed. Using the mAb grouped according to their binding specificity to NKH1, three different epitopes on the antigen were identified: NKH1a defined by antibodies, N3 (Leu-19) and N21 (FP2-11.14); NKH1b by N9 (N901) and N77 (NKH1A); and NKH1c by N63 (T-199) and N126 (L185) [Schubert *et al.*, N16.1]. In contrast, using rosetting techniques with chromium-chloride-treated erythrocytes, two groups of epitopes were described by Pietsch and Hadam (N16.2). They grouped antibodies N3 (Leu-19), N21 (FP2-11.14), N63 (T-199), and N126 (L185) as completely blocking rosetting of mAb T-199. N9 (N901) and N77 (NKH1A) only partially blocked T-199. Data from both groups provided evidence that antibodies N9 (N901) and N77 (NKH1A) recognize the same epitope on the NKH1 antigen. The differences observed by the two groups may be due to different methods used. Whereas our group tested four of the NKH1

van den Hart,

M. K., and

ersons, S. P.,  
307 (1987).  
., and Rosse,

hikawa, M.,  
Daha, M. In  
Blood Trans-  
fusion Society,  
y, Manches-

G., Martin,  
(1987).

, D. J., Getty,  
cinski, M. L.

, 13 (1989).

## 700 NK- and Non-lineage antigens — papers

CD56 N16




Fig. 1. Mean reactivity profiles of CD56 mAb.

6 N16

N16 CD56

NK- and Non-lineage antigens — papers 701

Table 1. CD56 antibodies

Workshop no.	mAb name	Submitter	Species*	Isotype	Com†		Wt‡ P§	Molecular weight (kDa)¶		Subm. spec.
					R	H		Submitter	Workshop	
N003	Leu-19	Lanier	M	Ig1	k	No	No	155	220/135	Unique
N009	N901	Griffin	M	IgG1		No	No	200	135	Unique
N021	FP2-11.14	Poncellet	M	IgG2a		Yes	nd	No	220/135	Unique
N063	T-199	Felckert	M	IgG1	k	nd	nd	Yes	220/135	Unique
N077	NKH1A	Ritz	M	IgM		Yes	nd	200		Unique
N126	L185	Lanier	M	IgM	k	Yes	nd	155		Unique

\*M, mouse.

†Com, complement binding: R, rabbit, H, human; nd, not done; No, does not fix C; Yes, does fix C.

‡W, Western blotting.

§P, formalin fixation.

¶Molecular weight of multiple polypeptide chains are separated by a solidus (/).



antibodies as second reagents, Pietsch and Hadam used only N63 (T-199) as a second antibody.

Consistent with prior results [6], cultured human thymocytes acquire NKH1 expression and non-MHC restricted cytotoxicity [Heiken *et al.*, N16.4]. Increasing expression of NKH1 antigen was seen with duration of thymocyte culture and the expression of NKH1 correlated with activation of NK- and T-lymphocytes. By analysing stimulated thymocytes on a clonal level, the NKH1 antigen was detected on cytotoxic as well as on non-cytotoxic clones.

Several laboratories reported strong reactivity of N-CAM/NKH1 panel antibodies with tumour cells derived from the neuroectoderm, such as neuroblastoma, medulloblastoma, retinoblastoma, pheochromocytoma, some melanomas, and small-cell lung carcinomas [Feickert *et al.*, N16.3]. Niedecken described N-CAM/NKH1 antibodies as also reacting with peripheral autonomic nerve fibres. A cDNA has been cloned for human N-CAM [10]. Unfortunately, only one mAb, Leu-19, was tested on transfectants, and was found to be positive.

The role of the N-CAM/NKH1 molecule in NK-cell function or activation could not be elucidated during this Workshop. Various laboratories tested the new antibody panel for induction or blocking of cytotoxic activity, or proliferation of NK-cells. None of the antibodies with specificity to NKH1 could significantly affect NK-cell function.

In summary, the CD56 cluster defined by six mAb from five different laboratories could be established during this Workshop. This could be based upon their identical phenotypic reactivity pattern, and molecular weight.

## Acknowledgements

This work was supported by DFG Schm 596/3-1.

## References

1. Ritz, J., Schmidt, R. E., Michon, J., Hercend, T., and Schlossman, S. F. *Adv. Immunol.* 42, 181 (1988).
2. Hercend, T., Griffin, J. D., Bensussan, A., Schmidt, R. E., Edson, M. A., Brennan, A., Murray, C., Daley, J. F., Schlossman, S. F., and Ritz, J. *J. clin. Invest.* 75, 932 (1985).
3. Griffin, J. D., Hercend, T., Beveridge, R., and Schlossman, S. F. *J. Immunol.* 130, 2947 (1983).
4. Lanier, L. L., Le, A. M., Civin, C. I., Loken, M. R., and Phillips, J. H. *J. Immunol.* 136, 4480 (1986).
5. Schmidt, R. E., Michon, J. M., Woronicz, J., Schlossman, S. F., Reinherz, E. L., and Ritz, J. *J. clin. Invest.* 79, 305 (1987).
6. Lanier, L. L., Le, A. M., Ding, A., Evans, E. L., Krensky, A. M., Clayberger, C., and Phillips, J. H. *J. Immunol.* 138, 2019 (1987).
7. Lanier, L. L., Tesd, R., Bindl, J., and Phillips, J. M. *J. exp. Med.* 169, 2233 (1989).
8. Cunningham, B. A., Hemperly, J. J., Murray, B. A., Prediger, E. A., Btackenbury, R., and Edelman, G. M. *Science* 236, 799 (1987).
9. Hemperly, J. J., Edelman, G. M., and Cunningham, B. A. *Proc. nat. Acad. Sci. USA* 83, 9822 (1986).
10. Barton, C. H., Dickson, G., Gower, H. J., Rowett, L. H., Putt, W., Elsom, V., Moore, S. E., Goridis, C., and Walsh, F. S., *Development* 104, 165 (1988).
11. Walsh, F. S. *Neurochem. Internat.* (in press).
12. Nguyen, C., Mattei, M. G., Mattei, J. M., Santoni, M. J., Goridis, C., and Jordan, B. R. *J. Cell Biol.* 102, 711 (1986).
13. Owens, G. C., Edelman, G. M., and Cunningham, B. A. *Proc. nat. Acad. Sci., USA* 84, 294 (1987).
14. Edelman, G. M. *Science* 219, 450 (1983).

## N16.1 Heterogeneity of the NKH1 molecule

JÖRG SCHUBERT, PETER UCIECHOWSKI, PATRICIA DELANY,  
KATHRIN WORDELMANN, and REINHOLD E. SCHMIDT

Six mAb from the New Antibody Panel have been identified as recognizing the NKH1 antigen according to their phenotypic pattern: N3 (Leu-19), N9 (N901), N21 (FP2-11.14), N63 (T-199), N77 (NKH1A), and N126 (L185). They all bind to various preparations of NK-cells

and to several tumour cell lines of myeloid and neuroectodermal origin [Schubert *et al.*, N16].

We tested the available Workshop antibodies with NKH1 specificity for functional effects on cloned human NK-cell lines. None of the NKH1 antibodies could